

Carlos Humberto Balcenas

Prepa Tec
Calculus I 3rd partial
Quiz # 1A

76

Name Carlos Humberto Balcenas Gonzalez Mat. AC1130142

- I. Determine if true or false for each of the following statements (10 points each)
- ~~F~~ The second derivative of $y = 2e^{x^3}$ is $\frac{d^2y}{dx^2} = 6xe^{x^3}(2+3x^3)$
 - ~~F~~ The derivative of $6x - 4x^2y = 2y^2 + 1$ is $\frac{dy}{dx} = \frac{-6}{8x+4y}$
 - ~~F~~ The derivative of $y = x^{2x}$ is $y' = 2x^{2x}(\ln(x)+1)$
 - ~~F~~ A spherical balloon is being inflated with a gas at a rate of 6 cm³ per second. Then the rate at which its radius is changing when its radius measures 8 cm is $\frac{dr}{dt} = \frac{3\pi}{128} \frac{cm}{sec}$. (Hint: $V = \frac{4}{3}\pi r^3$)

II. Answer the following problem. (10 points each letter)
A dynamite charge blows a rock up with a velocity of 160 ft/s. The height of the rock is given by the function $h(t) = 160t - 16t^2$ where "h" is measured in feet and "t" in seconds. Find the following:

- The equation that gives the velocity of the rock at any time.
 $v(t) = 160 - 32t$
- The time when velocity is zero (that is the time to reach the maximum height)
 $0 = 160 - 32t \implies -160 = -32t \implies \frac{-160}{-32} = \frac{-32t}{-32} \implies t = 5s$
- The maximum height of the rock (that is when velocity is zero)
 $h(t) = 160(5) - 16(5)^2$
 $h(t) = 800 - 400$
 $h(t) = 400 ft$
- The times (on the way up and on the way down) when the height is at 256 feet.
 $256 = 160t - 16t^2$
 $16t^2 - 160t + 256 = 0$
 $10 \pm \sqrt{10^2 - 4(1)(16)} = \frac{10 \pm \sqrt{100 - 64}}{2} = \frac{10 \pm \sqrt{36}}{2} = \frac{10 \pm 6}{2}$
 $t = \frac{10+6}{2} = 8$ and $t = \frac{10-6}{2} = 2$
23s and -13s
- The velocities of the rock when the height is 256 feet.
 $160 - 32(8) = -576$
 $160 - 32(2) = 576$
 $v = 160 - 32(2) \implies v = 160 - 64 \implies v = 96$
 $v = -576$
23s, 28s
- The equation that gives the acceleration of the rock at any time.
 $a(t) = -32$

Prepa Tec



Calculus I 3rd partial 65 Quiz # 2A

Name Carlos Humberto Balcenas Gonzalez Mat. AC1130142

- Determine if true or false for each of the following statements. (5 points each)
- ~~T~~ The function $f(x) = x^3 - 4$ has a relative maximum at $(0, -4)$
 - ~~T~~ If "a" is a critical point of the function $f(x)$ that is continuous, and if $f'(x) < 0$ at $(-\infty, a)$ and $f'(x) > 0$ at (a, ∞) , then, $f(x)$ has a relative minimum at $(a, f(a))$.
 - ~~T~~ Let f be a function whose second derivative exists on an open interval, if $f''(x) > 0$ for all x in that interval, then the graph of f is concave upward on that interval.
 - ~~F~~ The function $f(x) = -3x^3 - 3x^2 + 14$ has only one critical point.

- Choose the right answer (10 points each)
- ~~A~~ If $(c, f(c))$ is a critical point, then
A) $f'(c) = 0$ B) $f'(c) < 0$ C) $f'(c) > 0$ **D) $f'(c) = 0$**
 - ~~C~~ According to the second derivative test if $f''(c) > 0$, then
A) $f(c)$ is concave downward.
B) $f(c)$ is relative maximum.
C) $f(c)$ is a critical point.
D) $f(c)$ is a relative minimum.
 - ~~P~~ The function $f(x) = 20x - x^2$ has a critical point:
A) $x = 10$ B) $x = -10$ C) $x = 0$ D) $x = 1$
 - ~~A~~ If $(f''(x) > 0)$ then $f(x)$ is:
A) concave upward B) concave downward C) decreasing D) increasing
 - ~~B~~ It can be determined if the curve of $y=f(x)$ has a change of concavity:
A) Critical point B) Inflection point C) x-intersect D) y-intercept
 - ~~C~~ The function $y = -x^3 + 6x^2$ has a relative minimum at:
A) (4, 32) B) (0, 0) C) (4, 32) D) (6, 0)
 - ~~C~~ The function $y = x^3 - 3x^2$ has a relative maximum at:
A) (1, -2) B) (3, 0) C) (0, 0) D) (2, -4)

Answer the following showing your entire procedure.

1. The following graph represents $f(x)$ use it to sketch the graphs of $f'(x)$. (10 points)

$y = x^3 - 3x^2$
 $y' = 3x^2 - 6x$
 $3x(x-2)$
 $3x = 0 \implies x = 0$
 $x - 2 = 0 \implies x = 2$
 $(0)^3 - 3(0)^2 = 0 - 0 = 0$
 $(2)^3 - 3(2)^2 = 8 - 12 = -4$
 $(10, 0)$
 $8 - 12$

Name: Carla Humberto Bovera ID: A119414 Date: 30/01/2013

Consider each of the following situations and answer clearly. Remember to use the appropriate mathematical notation and to frame your final answer.

1. An object is moving along a straight line, and its position (in meters) is given by the function $s(t) = 80t - t^2$. Determine
- The velocity of the object when $t = 2$ sec. $v = 16 \text{ m/s}$
 - The acceleration when $t = 3$ sec. $a = -2 \text{ m/s}^2$
 - The time when the velocity is zero and the position of the object at that time. $t = 4 \text{ s}$, $s = 160 \text{ m}$

2. An object is moving along a straight line, and its position (in meters) is given by the function $s(t) = 3t + \frac{48}{t+1}$.

- Determine
- The velocity of the object when $t = 2$ sec. $v(t) = 3$
 - The acceleration when $t = 2$ sec. $a = 5 \text{ m/s}^2$
 - The time when the velocity is zero and the position of the object at that time. $t = 5 \text{ s}$, $s = 11 \text{ m}$

3. A dynamite charge blows a rock up with a velocity of 160 feet/sec. The height of the rock is given by $h(t) = 160t - 16t^2$

where "h" is measured in feet and "t" in seconds. Find

- The equation that gives the velocity of the rock at any time. $v = 160 - 32t$
- The time when the velocity is zero. $t = 5 \text{ s}$
- The height of the rock when the velocity is zero (maximum height). 400 m
- The times (on the way up and on the way down) when the height is 256 feet. 2 s , 4 s
- The velocities of the rock when the height is 256 feet. 46 m/s , -46 m/s
- The equation that gives the acceleration of the rock at any time. $a = -32$
- How long does it take the rock to fall back down? 10 s in one ideal

4. A baseball is thrown upward while being in the moon (hypothetically), with an initial velocity of 24 meters/second. The height of the ball is given by $s = 24t - 0.8t^2$

- Find the equations of velocity and acceleration at any time. $v = 24 - 1.6t$, $a = -1.6$
- How long does it take the ball to reach its maximum height? 1.5 s
- Find the maximum height of the ball. 180 m
- How long was the ball in the air? 30 s in one ideal

5. The position of an object is given by $S(t) = t^3 - 6t^2 + 9t$ where "t" is measured in seconds and "s" in meters.


- Find the equations of velocity and acceleration as a function of time. $v = 3t^2 - 12t + 9$, $a = 6t - 12$
- Find the time when the velocity is zero. $t = 3 \text{ s}$
- Find the acceleration when the velocity is zero. $a = 6 \text{ m/s}^2$
- Find the time when the acceleration is zero and then give the velocity at that time. $t = 2 \text{ s}$, $v = 9 \text{ m/s}$

6. The height of a certain tree (starting from being 1 year old) is modeled by $H(t) = 5\sqrt{t} + 2t^2 + 10$, where height is measured in cm and time in years

Find:

- The height of the tree in its 5th year (hint $t=4$). $h(5) = 82 \text{ cm}$
- The function that models the rate of change of its height. $h'(t) = \frac{5}{2\sqrt{t}} + 4t$
- The rate of change when $t=4$. 31 cm
- The rate of change when $t=9$. 36 cm
- When is the tree growing faster? at $t=4$ or $t=9$ years? Why? $t=9$ because it grows more than tree

CHALLENGE: The following graph shows the position of a particle that moves along a straight line (author: Lic. Norma Patricia Salinas Martínez).



- In which interval or intervals is the velocity of the particle positive?
- In which interval or intervals is the velocity of the particle negative?
- In which interval or intervals of time is the position increasing slower?
- In which interval or intervals of time is the position increasing faster?
- In which interval or intervals of time is the position decreasing slower?
- In which interval or intervals of time is the position decreasing faster?
- In which interval or intervals of time is the velocity increasing?
- In which interval or intervals of time is the velocity decreasing?

Carla Humberto Bovera ID: A119414

1. Find the derivative of given functions, using logarithmic differentiation

1) $y = x^{2x}$, $x > 0$

$$\ln y = \ln(x^{2x})$$

$$\ln y = 2x \ln x$$

$$\ln y = 2x \ln x + \frac{2x}{x} \ln x$$

$$y' = \frac{2x}{y} y' + \frac{2x}{x} \ln x + \frac{2x}{x} \ln x$$

3) $y = 4x\sqrt{x^2+1}$, $x > 0$

$$\ln y = \ln(4x\sqrt{x^2+1})$$

$$\ln y = \ln 4x + \ln(x^2+1)^{1/2}$$

$$\ln y = \ln 4x + \frac{1}{2} \ln(x^2+1)$$

$$\ln y = \frac{2x}{y} y' + \frac{2x}{2x^2+1} \ln(x^2+1)$$

5) $y = \frac{x^2\sqrt{x-4}}{(x+3)^2}$, $x > 4$

$$\ln y = \ln \left(\frac{x^2\sqrt{x-4}}{(x+3)^2} \right)$$

$$\ln y = \ln(x^2\sqrt{x-4}) - \ln(x+3)^2$$

$$\ln y = 2 \ln x + \frac{1}{2} \ln(x-4) - 2 \ln(x+3)$$

7) $y = x^{\sqrt{x}}$, $x > 0$

$$\frac{x \sqrt{x}}{2\sqrt{x}} (x)$$

9) $y = [\sin(x)]^{3x}$, $\sin(x) > 0$

$$y' = 3x \cos x \times 3x$$

2) $y = x^{e^x}$, $x > 0$

$$\ln y = \ln(x^{e^x})$$

$$\ln y = e^x \ln x$$

$$\ln y = e^x \ln x + \frac{e^x}{x} \ln x$$

4) $y = \sqrt{\frac{x^2-4}{x^2+1}}$, $x > 2$

$$\ln y = \ln \left(\frac{x^2-4}{x^2+1} \right)^{1/2}$$

$$\ln y = \frac{1}{2} \ln \left(\frac{x^2-4}{x^2+1} \right)$$

6) $y = \frac{(3+4x)^3}{(1-2x)^3}$, $-\frac{3}{4} < x < \frac{1}{2}$

$$\ln y = \ln \left(\frac{(3+4x)^3}{(1-2x)^3} \right)$$

$$\ln y = 3 \ln(3+4x) - 3 \ln(1-2x)$$

8) $y = x^{\sin(x)}$, $x > 0$

$$y' = \frac{(\sin x)^x + \cos x \ln x \cdot x^{\sin x}}{x}$$

10) $y = -x^{\sin(x)}$, $x > 0$

$$y' = \left(\frac{-\sin x + \cos x \ln x \cdot x}{x} \right) \times \sin x \cdot x$$



By: Arq. Monica M. Paniagua, Ing. Ziad Najjar, Lic. Lucy Solis, Lic. Carmela de la fuente

Name: Carla Humberto Bovera ID: A119414 Date: 30/01/2013

Related rates

1. A spherical balloon is being filled at a rate of $50 \text{ in}^3/\text{sec}$, at what rate does the radius increase when the radius is 5 in?

$$\frac{dV}{dt} = 50 \text{ in}^3/\text{sec}$$

$$V = \frac{4}{3} \pi r^3$$

$$\frac{dV}{dt} = 4\pi r^2 \frac{dr}{dt}$$

$$50 = 4\pi (5)^2 \frac{dr}{dt}$$

$$\frac{dr}{dt} = \frac{50}{100\pi} = \frac{1}{2\pi} \text{ in/sec}$$

2. The area of a circle is increasing at a rate of $20 \text{ in}^2/\text{min}$. Find the rate at which the radius is increasing when the radius is 4 in.

$$\frac{dA}{dt} = 20 \text{ in}^2/\text{min}$$

$$A = \pi r^2$$

$$\frac{dA}{dt} = 2\pi r \frac{dr}{dt}$$

$$20 = 2\pi (4) \frac{dr}{dt}$$

$$\frac{dr}{dt} = \frac{20}{8\pi} = \frac{5}{2\pi} \text{ in/min}$$

3. A stone is thrown into a lake and a circular ripple moves out at a constant rate of 0.5 meters/sec. Find the rate at which the circle's area is increasing at $r = 0.4$ meters.

$$\frac{dA}{dt} = 2\pi r \frac{dr}{dt}$$

$$\frac{dA}{dt} = 2\pi (0.4)(0.5)$$

$$\frac{dA}{dt} = 0.4\pi \text{ m}^2/\text{s}$$

4. Air is being pumped into a spherical balloon making the radius change at a constant rate of 0.5 cm/sec. Find the rate of change of the volume and the rate of change of the surface area when the radius is 10 cm ($V = \frac{4}{3}\pi r^3$, $A = 4\pi r^2$)

$$\frac{dV}{dt} = 4\pi r^2 \frac{dr}{dt}$$

$$\frac{dV}{dt} = 4\pi (10)^2 (0.5)$$

$$\frac{dV}{dt} = 200\pi \text{ cm}^3/\text{s}$$

$$\frac{dA}{dt} = 8\pi r \frac{dr}{dt}$$

$$\frac{dA}{dt} = 8\pi (10) (0.5)$$

$$\frac{dA}{dt} = 40\pi \text{ cm}^2/\text{s}$$

5. A cone is increasing in size as time goes by in such a way that the volume is changing at a constant rate of $75 \text{ cm}^3/\text{min}$. The height is twice the radius. Determine the rate of change of the height, when the height is 5 cm. ($V = \frac{1}{3}\pi r^2 h$)

$$\frac{dV}{dt} = 75 \text{ cm}^3/\text{min}$$

$$V = \frac{1}{3}\pi r^2 h$$

$$\frac{dV}{dt} = \pi r^2 \frac{dh}{dt} + \frac{2}{3}\pi r h \frac{dr}{dt}$$

$$75 = \pi (2.5)^2 \frac{dh}{dt} + \frac{2}{3}\pi (2.5)(5) \frac{dr}{dt}$$

$$75 = 6.25\pi \frac{dh}{dt} + 25\pi \frac{dr}{dt}$$

$$\frac{dh}{dt} = \frac{75 - 25\pi \frac{dr}{dt}}{6.25\pi}$$